

***Enicmus adrianae* sp. nov. – a new scavenger beetle (Coleoptera: Latridiidae) from Baltic amber**

Andris Bukejs, Hans-Peter Reike, Wolfgang H. Rucker

Bukejs A., Reike H.-P., Rucker W.H. 2012. *Enicmus adrianae* sp. nov. – a new scavenger beetle (Coleoptera: Latridiidae) from Baltic amber. *Baltic J. Coleopterol.*, 12(2): 149 - 154.

Enicmus adrianae sp. nov. (Coleoptera: Latridiidae) from Baltic amber is described and illustrated in the current paper. It is the first described Eocene species of the genus.

Key words: Coleoptera, Latridiidae, *Enicmus adrianae*, new species, Baltic amber, Upper Eocene.

Andris Bukejs. Institute of Systematic Biology, Daugavpils University, Vientbas 13, Daugavpils, LV-5401, Latvia. E-mail: carabidae@inbox.lv

Hans-Peter Reike. Waldemarstraße 8, 01139 Dresden, Germany. E-mail: latridiidae@gmx.de

Wolfgang H. Rucker. Von-Ebner-Eschenbach-Str. 12, D-56567 Neuwied, Germany. E-mail: Coleoptera@Latridiidae.de

INTRODUCTION

Latridiidae Erichson, 1842 is a medium sized family of Coleoptera with approximately 800 described species worldwide. There are ca. 170 scavenger beetle species known in Europe (Rucker 2011). The tropical rainforests of Africa and South America are the biggest diversity centers for some genera (*Melanophthalma* Motschulsky, 1866 and other) of the family Latridiidae (Rucker 1981, 1984). The genus *Enicmus* C. G. Thomson, 1859 contains 50 species (Rucker 2010).

Only a few species of Latridiidae are known from fossils (Borowiec 1985; Bukejs et al. 2011; Hawkeswood et al. 2009; Heer 1856; Kirejtshuk & Azar 2008; Kirejtshuk & Nel 2009; Kirejtshuk et al. 2009; Schlechtendal 1894; Wickham 1913,

1914a, 1914b; Zherikhin 1977). Thereby the most indications contain not more than a generic attribution without any descriptions (Grimaldi & Engel 2005; Hieke & Pietrzeniuk 1984; Klebs 1910; Kubisz 2000, 2001; Poinar & Poinar 1999; Rasnitsyn & Ross 2000; Spahr 1981; etc). The detailed review of this family in the fossil records is given in the catalogue by Ponomarenko & Kirejtshuk (2012).

In the current paper a new species of the genus *Enicmus* C.G. Thomson, 1859 is described from Baltic amber, which is usually dated as the Upper Eocene, although sometimes older (even the Lowermost Eocene) (Weitschat & Wichard 2010). It is the first described fossil species of the genus.

MATERIAL AND METHODS

The material examined is deposited in the collection of the Museum of Natural History, Institute of Systematics and Evolution of Animals, Polish Academy of Sciences (Krakow, Poland), in the collection of PAW Museum of the Earth (Warszawa, Poland), and in the private collection of Wolfgang Rucker (Neuwied, Germany).

The photos were taken using a stereomicroscope Nikon SMZ 745T and digital sight camera Nikon DS-F11. Merging of layers was done with CombineZP.

SYSTEMATIC PALEONTOLOGY

Latridiidae Erichson, 1842

Latridiinae Erichson, 1842

***Enicmus* C.G. Thomson, 1859**

***Enicmus adrianae* sp. nov.**

Figs 1-2.

Material

Holotype – “MP/564”, sex unknown, Baltic Amber; deposited in the collection of the Museum of Natural History, Institute of Systematics and Evolution of Animals, Polish Academy of Sciences, Krakow, Poland. A rather well visible complete beetle with indistinctly visible mouthparts and base of antennae. The specimen is included in a small and thin elongate amber piece (with length about 20 mm and width 6 mm, weight 0.2 g).

Paratype – No. 0028/2001, sex unknown, Baltic amber, Yantarny village, Kaliningrad region, Russia; deposited in the collection of Wolfgang Rucker (Neuwied, Germany).

Type strata

Baltic Amber, Upper Eocene.

Etymology

The epithet of the new species is devoted to the first authors' daughter, Adriana Bukeja.

Diagnosis

Enicmus adrianae sp. nov. is most closely related to the recent *E. amici* Lohse, 1981, *E. histrio* Joy & Tomlin, 1910 and *E. transversus* (Olivier, 1790), which have (just like the new species) elytral interspaces smaller than a puncture in diameter and the humeral row of the elytra is convex in the posterior half. However, the new species differs from them in the shape of the first antennomere and the more dense and deep puncture of the first abdominal ventrite.

Some other species of the genus, resp. *Enicmus planipennis* Strand, 1940, *E. fungicola* C. G. Thomson, 1868, *E. testaceus* (Stephens, 1830) and *E. rugosus* (Herbst, 1793) have interspaces between the elytral rows double as wide as the puncture, but in *E. adrianae* sp. nov. elytral interspaces are smaller than a puncture in diameter. The humeral row of the elytrae is slightly convex in the posterior half of the elytrae in *E. adrianae* sp. nov., but in *E. planipennis*, *E. testaceus* and *E. rugosus* it is convex in the anterior half of the elytrae, in *E. fungicola* convex in the posterior half of the elytrae. Further the shape of the scape and the shape of the pronotum differs between the species.

In *E. atriceps* Hansen, 1962 just like in *E. adrianae* sp. nov. elytral interspaces are smaller than a puncture in diameter (except the second interspace), but the shape of the scape and the shape of the pronotum differs between the species.

There are no described fossil species of the genus *Enicmus*. From the other 4 fossil species of the subfamily Latridiinae (*Archelatrius marinae* Kirejtshuk & Azar, 2009, *Latridius kulickae* Borowiec, 1985, *L. jantaricus* Borowiec, 1985, *L. alexeevi* Bukejs, Kirejtshuk, Rucker 2011) *Enicmus adrianae* sp. nov. clearly differs in convex (but not carinate) elytral interstriae, antennal structure and some other characters.

Description

Holotype. Length 1.5 mm, max. width 0.59 mm. Dorsal surface brown, glabrous; legs and anten-

nae rufous. Elongate, strongly convex dorsally and moderately convex ventrally.

Head transverse, evenly convex dorsally, covered with dense and moderately large punctures markedly larger than eye facets in diameter, interspaces between them smooth and distinctly less than a puncture diameter. Head with broad and distinct median sulcus at its base. Eyes large, convex with moderately large and distinct facets; distance between them about 5 times as long as width of one eye. Tempora are not larger than the width of one eye. Neck markedly separated. Anterior part of head and mouthparts invisible because of milky cover. Antennal insertions located slightly before anterior edge of eyes. Antennae 11-segmented, moderately long, reaching posterior 1/3 of pronotum. Scape is not distinctly visible, subtriangular, larger than antennomere 2; antennomere 2 larger than antennomere 3; antennomeres 3-8 slightly (1.3-1.7 times) longer than wide, widened apically. Antennal club loose, distinctly 3-segmented; antennomere 11 widely suboval and acute apically; antennomeres 9-10 subequally large, slightly slender and about 1.8 times shorter than ultimate antennomere.

Pronotum transverse, approximately 1.2 times wide as long; slightly narrowed anterior and more posterior; widest in anterior 1/3; lateral margins in anterior half rounded and in basal 1/3 subparallel; covered with dense and moderately large punctures (as on the head). Anterior angles slightly acute, weakly prominent anteriorly; posterior angles subrectangular, weakly rounded. Disc convex and moderately steeply sloping to sides. Basal, anterior and lateral sides completely bordered. Pronotum with transverse impression at the base, oval impression near each posterior angle, and two longitudinal subelongate impressions on the disc. Scutellum small, subtriangular.

Elytra broadly oval, combined about 1.5 times as long as wide, widest behind the middle; strongly convex at disc and rather steeply sloping to sides, lateral sides slightly explanate. Base of elytra distinctly wider than the base of the pronotum. Hu-

meri distinct. Punctures in elytral rows dense, large, becoming smaller in posterior half, in basal 1/3 of the elytra punctures distinctly larger than punctures on the pronotum; interspaces between punctures smaller than a puncture diameter (especially in the basal half). Punctures are forming 8 regular rows. All interstriae smooth, convex, interstriae 6, 7 and 8 stronger convex than other interstriae.


Pro-, meso- and metathorax covered with moderately dense and large punctures (approximately as punctures on the head), interspaces between punctures smaller than a puncture in diameter. Procoxae separated by a prosternal process; process narrow and prominent beyond posterior margin of procoxae. Procoxal cavities broadly closed posteriorly. Pro- and mesocoxae suboval, metacoxae transversely oval. Distance between mesocoxae distinctly greater than that between procoxae; and between metacoxae about twice as great as that between mesocoxae. Metasternum flattened in the middle. Epipleura of elytra wide and weakly narrowed in apical part; with distinct punctures. The first abdominal ventrite is the longest one, with dense and moderately large punctures (comparable with the puncture on the thorax); ventrites 2-5 covered with finer and sparser punctures.

Legs moderately long and narrow. Trochanters slightly elongate. Femora spindle-shaped, thickest at the middle and about 2-3 times as wide as tibiae. Tibiae slender, tibiae and femora approximately comparable in length. Tarsi with three simple subcylindrical tarsomeres, about 2/3 as long as tibiae; tarsomeres 1 and 2 approximately comparable in length; tarsomere 3 markedly, about 1.5 times longer than both previous ones combined. Claws simple, small and thin.

Variation

Body size of paratype – length 1.6 mm, width 0.63 mm.

NOMENCLATURAL NOTE



Figs 1-2. *Enicmus adrianae* sp. nov., holotype: 1 – habitus, latero-dorsal view, 2 – habitus, latero-ventral view.

Latridius jantarius Borowiec, 1985 was placed into the genus *Stephostethus* Le Conte, 1878 by Kubisz (2000). According to the characters of the holotype (deposited in the collection of PAW Museum of the Earth) of this species it belongs to the genus *Latridius* Herbst, 1793 and it is necessary to replace this species back – *Latridius jantarius* Borowiec, 1985 stat. rev.

ACKNOWLEDGEMENTS

The authors are sincerely grateful to Aneta Garbula (Museum of Natural History, Institute of Systematics and Evolution of Animals, Polish Academy of Sciences, Krakow, Poland) for the loan of the material. Special thanks go to Prof. Dr. hab. Barbara Kosmowska-Ceranowicz (PAW Museum of the Earth, Warszawa, Poland) as well as to Dr. Wolfgang Weitschat (Geologisches

Institut Universität Hamburg, Hamburg, Germany) and to Carsten Gröhn (Glinde, Germany) for the realization of the loan of the holotype of *L. jantarius*. The authors are also grateful to Dr. Alexander G. Kirejtshuk (Zoological Institute of the Russian Academy of Sciences, St. Petersburg, Russia) for constructive advice and valuable comments on our manuscript.

REFERENCES

- Borowiec L. 1985. Two new species of *Latridius* sensu lato (Coleoptera: Latridiidae) from Baltic Amber. *Polish Journal of Entomology*, 55: 251–254.
- Bukejs A., Kirejtshuk A.G., Rucker W.H. 2011. New species of *Latridius* (Coleoptera: Latridiidae) from Baltic amber. *Baltic Journal of Coleopterology*, 11 (2): 203–207.

- Grimaldi D., Engel M. S. 2005. *Evolution of the Insects*. Cambridge University Press, New York: 1-755.
- Hawkeswood T.J., Makhan D., Turner J.R., 2009. *Latridius jacquelinae* sp. nov., a new scavenger beetle from Baltic amber (Coleoptera: Latridiidae). *Entomologische Zeitschrift*, Stuttgart, 119(3): 108-110.
- Heer O. 1856. Über die fossilen Insekten von Aix in der Provence. *Vierteljahrsschrift der Naturforschenden Gesellschaft in Zürich* 1: 1-40, pl. 1, 2.
- Hieke F., Pietrzeniuk E. 1984. Die Bernstein-Käfer des Museums für Naturkunde, Berlin (Insecta: Coleoptera). *Mitteilungen aus dem Zoologischen Museum Berlin*, 60 (2): 297-326.
- Kirejtshuk A., Nel A. 2009. New genera and species of Cucujiformia (Coleoptera, Polyphaga) from lowermost Eocene French amber. *Denisia* 26, zugleich *Kataloge der Oberösterreichischen Landesmuseen Neue Serie*, 86: 103-118.
- Kirejtshuk A.G., Azar D. 2008. New taxa of beetles (Insecta, Coleoptera) from Lebanese amber with evolutionary and systematic comments. *Alavesia*, 2: 15-46.
- Kirejtshuk A.G., Azar D., Tafforeau P., Boistel R., Fernandez V. 2009. New beetles of Polyphaga (Coleoptera, Polyphaga) from Lower Cretaceous Lebanese amber. *Denisia*, 26, zugleich *Katalog der oberösterreichischen Landesmuseen, Neue Serie*, 86: 119-130.
- Klebs R. 1910. Über Bernsteineinschlüsse in allgemeinen und die Coleopteren meiner Bernsteinsammlung. *Schriften der physikalisch-ökonomischen Gesellschaft zu Königsberg*, 51: 217-242.
- Kubisz D. 2000. Fossil beetles (Coleoptera) from Baltic amber in the collection of the Museum of Natural History of ISEA in Kraków. *Polish Journal of Entomology*, 69 (2): 225-230.
- Kubisz D. 2001. Beetles in the collection of the museum of amber inclusions, University of Gdańsk, with description of *Colotes sambicus* sp. n. (Coleoptera: Melyridae). *Polish Journal of Entomology*, 70 (4): 259-265.
- Poinar G.O. Jr., Poinar R. 1999. *The Amber Forest: A reconstruction of a Vanished World*. Princeton University Press, Princeton & Oxford: xviii, 239 pp.
- Ponomarenko A.G., Kirejtshuk A.G. 2012. Catalogue of fossil Coleoptera. Available at <http://www.zin.ru/Animalia/Coleoptera/rus/paleosys.htm> [accessed January 2012]
- Rasnitzyn A.P., Ross A.J. 2000. A preliminary list of arthropod families present in the Burmese amber collection at the Natural History Museum, London. *Bulletin of the Natural History Museum, Geological Series*, 56 (1): 21-24.
- Rücker, W. H. 1981. Revision der Motschulsky-Typen aus Nord- und Südamerika, Coleoptera: Lathridiidae, *Melanophthalma* Motschulsky. *Entomologische Blätter*, 77 (1-2): 49-53.
- Rücker, W. H. 1984. Neue Arten der Gattung *Melanophthalma* Motschulsky aus Afrika (Coleoptera, Lathridiidae). *Annales Historico-Naturales Musei Naturalis Hungarici*, 76: 139-141
- Rücker W. H. 2010. Checklist Latridiidae & Merophysiinae of the World. *Latridiidae & Merophysiinae*, 9: 1-16.
- Rücker W. H. 2011. Fauna Europaea: Coleoptera, Latridiidae. In Audisio P. (ed.) *Fauna*

Europaea: Coleoptera, Beetles, Fauna
Europaea version 2.4, <http://www.faunaeur.org>. [accessed March 2012]

Received: 12.06.2012.

Accepted: 15.12.2012.

- Schlechtendal, D. H. R. von 1894. Beiträge zur Kenntnis fossiler Insekten aus dem Braunkohlengebirge von Rott am Siebengebirge. *Abhandlungen der Naturforschenden Gesellschaft zu Halle*, 20: 197-228, plates 12–14.
- Spahr U. 1981. Systematischer Katalog der Bernstein- und Kopal-Käfer (Coleoptera). *Stuttgarter Beiträge zur Naturkunde, Ser. B* 80: 1-107.
- Weitschat W., Wichard W. 2010. Baltic amber. In: Penney D. (ed.): *Biodiversity of fossils in amber from the major world deposits*. Siri Scientific Press, Manchester: 80-115.
- Wickham H.F. 1913. Fossil Coleoptera from Florissant in the United States National Museum. *Proceedings of the United States National Museum*, 45: 283-303.
- Wickham H.F. 1914a. Twenty new Coleoptera from the Florissant shales. *Transactions of the American Entomological Society*, 40: 257-270, 8 pls.
- Wickham H.F. 1914b. New Miocene Coleoptera from Florissant. *Bulletin of the Museum of Comparative Zoology at Harvard College*, 58 (11): 423-494, 16 pls.
- Zherikhin V.V. 1977. Infraorder Cucujiformia. Family Acanthocnemidae. Family Lathridiidae. In: Arnoldi L.V., Zherikhin V.V., Nikritin L.M., Ponomarenko A.G. *Mesozoiskie zhestkokrylye [Mesozoic beetles]*. *Trudy Paleontologicheskogo Instituta AN SSSR [Proceedings of the Palaeontological Institute of the USSR Academy of Sciences]*, 161. Moscow, Nauka: 135-142. [in Russian]